

A Review of Perineal Talc Exposure and Risk of Ovarian Cancer

Bernard L. Harlow, Ph.D.¹ and Patricia A. Hartge, Sc.D.²

¹Obstetrics and Gynecology Epidemiology Center, Brigham and Women's Hospital, Harvard Medical School

²Division of Cancer Etiology, National Cancer Institute, National Institutes of Health

Running Title: Perineal Talc Exposure and Risk of Ovarian Cancer

Address Correspondence to:

Bernard L. Harlow, Ph.D.
Obstetrics and Gynecology Epidemiology Center
Brigham and Women's Hospital
221 Longwood Avenue (BLI-3)
Boston, MA 02115

The authors provide a detailed review of the events that led to the interest in talc as a possible ovarian carcinogen, the epidemiological studies published to date, and their perspective on the interpretation of the findings including potential limitations, biases, and issues surrounding the plausibility of a causal association. The authors conclude that the range of relative risk estimates from epidemiology, 1.0 to 1.8, is plausible, but that additional epidemiologic studies, especially prospective investigations are needed. In addition, clinico-pathological studies are needed to confirm or deny the reports of talc embedded in human ovarian tissue and reports of talc migration through the human female reproductive tract.

Talc is a natural mineral, hydrous magnesium silicate, that is formed by the hydrothermal alteration of chrysotile and tremolite, two types of the asbestos group minerals (Cralley et al., 1968; Blejer et al., 1973). Because of its chemical similarity to asbestos, it has been implicated as both a lung (Thomas et al., 1987; Kleinfeld et al., 1974) and ovarian carcinogen (Longo and Young, 1979; Cramer et al., 1982; Harlow et al., 1992). Yet there is a surprisingly small body of literature to support its ovarian carcinogenicity, especially with respect to the pathological identification of embedded particulates within ovarian tissue (Henderson et al., 1971; Henderson et al., 1979). Although several controlled epidemiological studies have investigated the talc and ovarian cancer association, few have collected enough details on methods and frequency of exposures to make a meaningful assessment. Nevertheless, its notoriety has grown to the extent that a special workshop on the health consequences of talc exposure was recently sponsored by the Food and Drug Administration and the International Society of Regulatory Toxicology and Pharmacology. As part of our contribution to the understanding of perineal talc exposure and risk of ovarian cancer, we have summarized the earlier research that forewarned of a possible association, the results from more recent epidemiological studies, and offer our assessment regarding the plausibility of such an association in light of the potential limitations and biases inherent in case-control studies.

PATHOLOGICAL AND CLINICAL STUDIES

Many pathological and clinical observations preceded epidemiological investigations of talc and ovarian cancer. Interest in talc as an ovarian

carcinogen dates back more than 50 years when postoperative talc granulomas were first reported in the literature (Eiseman et al., 1947). Eiseman (1947) published a summary of more than 20 independent reports of talc induced granulomas following surgery that were reported in the literature between 1933 and 1947. These reports included results from clinical series of patients, laboratory-confirmed presence of talc crystals within granulomatous tissues, and inflammatory reactions as a consequence of subcutaneous and intraperitoneal injections of talc-suspensions in laboratory animals.

Using an extraction-replication technique developed to examine asbestos within the tissue of various mesotheliomas, Henderson (1971) examined multiple sections of ovarian tumors. No asbestos particles were found, but talc particulates were observed in 75% of 13 tumors examined (Henderson, et al., 1971). This finding was greeted with skepticism (Hildick-Smith, 1976), partly because of the possible contamination by talc on surgical gloves at the time of specimen collection and processing. A reevaluation of 9 new specimens collected from deep within tissue samples extracted with the use of forceps and noncontaminated surgical gloves reported the presence of talc particulates in all 9 specimens, including 3 from ovaries free of disease (Henderson, et al., 1979). Surprisingly, no further pathological analyses of talc particulates within ovarian tumor tissue have been reported in the literature.

ASBESTOS AND OVARIAN CANCER

Asbestos is the common name given to a number of naturally occurring hydrated mineral (inorganic) silicates. The four most common types are chrysotile, amosite, crocidolite and anthophyllite. Two other types, tremolite and actinolite, are found less commonly. Each of these inorganic silicates are

chemically different from each other (Michaels et al., 1979). By the mid-1970's, a body of evidence was mounting implicating asbestos exposure as a potential risk factor for ovarian cancer. The link between asbestos and ovarian cancer was first reported by Keal (1960) when it was noted that women suffering from asbestosis appeared to have concurrent ovarian neoplasms more frequently than the general population of other women at the London Hospital. During a five year retrospective search of hospital records, 9 of 23 women admitted with a diagnosis of asbestosis had been diagnosed with an intra-abdominal neoplasm as well. Shortly thereafter, Newhouse et al. (1985) identified 9 deaths associated with ovarian neoplasms (3.6 expected) from a cohort of 922 women first employed at an asbestos factory between 1933 and 1942 and followed for more than 38 years. Of these nine cases, 5 were in women who received severe exposure for more than two years in which only 0.9 cases should have been expected.

Three retrospective cohorts that totalled 1,827 women employed in the manufacturing of gas masks during world war II and followed for at least 30 years identified 23 deaths due to ovarian cancer compared to 10.6 deaths expected (Wignal et al., 1982; Acheson et al., 1982). The gas masks contained filter pads consisting of activated charcoal, merino wool and West Australian crocidolite or chrysotile. Although the intensity of the exposure is at best crudely estimated, the observed to expected ratio for developing a subsequent ovarian tumor appeared to be greater among women more severely exposed.

There are several factors that contribute to the biological plausibility of an asbestos-ovarian cancer association. First, Graham and Graham (1967) reported epithelial changes in the ovaries of guinea pigs and rabbits receiving intraperitoneal injections of asbestos products. Secondly, peritoneal

mesotheliomas closely resemble ovarian carcinoma histologically and if the tissue of origin is unknown may often be indistinguishable. Some believe they are in fact the same tumor since epithelial ovarian tumors are unlike epithelial tumors elsewhere in the body and because they arise from the "germinal epithelium" which is of mesodermal origin (Parmley and Woodruff, 1974). In fact, studies have reported that some women receiving prophylactic oophorectomy because of a strong family history of ovarian cancer have subsequently developed intra-abdominal lesions resembling ovarian tumors (Tobacman et al., 1982). Thirdly, environmental substances can penetrate the female peritoneal cavity through the vagina and fallopian tubes whereas no such entry is available in males (Egli and Newton, 1961; Henderson et al., 1986; Venter and Iturralde, 1979). This is consistent with the rarity with which peritoneal mesotheliomas occur in men.

EPIDEMIOLOGIC STUDIES

Since talc and asbestos are chemically similar and occur in nature together, there was some concern regarding the degree to which talc containing products were free of asbestiform fibers. A study of 21 consumer talcum powders labeled as baby powders, facial powders, or body powders obtained from retail stores in New York City between 1971 and 1975, reported that 10 contained concentrations of tremolite and anthophyllite ranging from 0.2 to 14 percent (Rohl et al., 1976) but these were not necessarily asbestiform. Although voluntary guidelines to limit the content of asbestiform fibers in consumer talcums were proposed by the cosmetic industry in 1976 (C.F.T.A., 1976), the magnitude of the risk of ovarian cancer as a result of prior perineal exposure to talc had never been quantitated. Because of this, a number of epidemiological studies during the late 1970's through the most

recent decade began to investigate talc as a potential risk factor for ovarian cancer.

A summary of the nine interview studies that have retrospectively ascertained perineal exposure to talc among women with and without ovarian neoplasms is presented in Table 1 (Cramer et al., 1982; Hartge et al., 1983; Whittemore et al., 1988; Harlow and Weiss, 1989; Booth et al., 1989; Harlow et al., 1992; Chen et al., 1992; Rosenblatt et al., 1992; Tzonou et al., 1993). All of the studies reported an overall estimate of ovarian cancer risk in women who reported any perineal exposure to talc compared to women with no such exposure. Risk estimates were greater than 1.0 in all but two of the studies. However, only two studies reported statistically significant odds ratios. Of the five studies that reported risk of ovarian cancer as a consequence of applying talc-containing powders to sanitary napkins, only one reported a significant association. Among the 7 studies that assessed talc exposure to diaphragms, 4 reported odds ratios greater than 1.0 and three reported odds ratios less than 1.0. None of them reached statistical significance. There were five studies that assessed exposure through direct perineal application after bathing. All five studies reported odds ratios greater than 1.0, but statistical significance was reported in only one study.

Three studies provided inconsistent results on whether frequency of perineal talc exposure, or years of exposure influenced ovarian cancer risk. Neither study by Whittemore (1988) or Booth (1989) showed a significant increase in risk with increasing applications per month, or years of talc exposure, whereas the study by Harlow et al (1992) showed a marginally significant trend of increasing ovarian cancer risk with increasing applications per month ($p=0.05$) and increasing years of exposure ($p=0.07$).

CONFIDENTIAL

Because the risk of any one outcome is dependent upon both the frequency and length of the exposure, Harlow et al (1992) created a continuous measure of total lifetime applications for each case and control. In Table 3, odds ratios are shown in columns of categories of lifetime applications including none, no more than 1,000, 1,000 to 10,000, and 10,000 or more applications. Although the odds ratios increased with increasing lifetime applications, the trend was marginally significant ($p=0.09$). After excluding talc exposure after tubal ligation or hysterectomy, little change in the odds ratios or test of trend was observed. However, when nonovulatory periods of exposure were excluded (i.e. exposure during pregnancy or breast feeding, while on oral contraceptives, or after menopause), there was a sizable increase in risk among those with more than 10,000 applications of talc which persisted after multivariate adjustment for parity, education, marital status, douching, weight, and use of sanitary napkins.

Two studies sought to determine whether the brand of talc-containing products influenced ovarian cancer risk. A study of borderline tumors by Harlow and Weiss (1988) reported that women who applied deodorizing powders with or without baby powder had nearly 3 times the risk of developing borderline ovarian tumor compared with women who reported no perineal use of powder ($RR=2.8$, $95\% \text{ C.I.}=1.1-11.7$), whereas women who applied only baby powder had no increase in risk of borderline tumors compared to nonusers. Attempts to replicate this finding in a different population of women with and without malignant ovarian tumors showed no such differences, mainly due to the lack of women reporting the use of deodorizing powders, and the likelihood of potential recall bias in obtaining an accurate exposure assessment (see following section).

CONFIDENTIAL

Lastly, in the study by Harlow et al, the authors stratified talc exposure by those whose use occurred only after 1960, and those who reported any talc use before 1960 with the idea that if the carcinogenic potential of talc is due to asbestos contamination, it is likely to manifest a greater risk in women who applied older talc-containing products. There was a 70% increase in risk among women who used talc before 1960 (95%CI=1.1-2.7) and only a 10% increase in risk (95%CI=0.6-2.1) among women whose use was exclusively after 1960, compared to nonexposed women.

LIMITATIONS AND BIASES IN RESULTS FROM EPIDEMIOLOGICAL STUDIES

The current data all derive from case-control studies, and their limitations stem from three primary sources: selection of controls that may not accurately reflect the underlying study base or population from which the cases derive, incomplete inclusion of the selected subjects because of refusals and other forms of nonresponse, and errors in the data collected, principally because of errors in recollection or reporting of talc exposures. To estimate the possible impact of these limitations, one can compare the nine studies to each other and use data from ancillary sources.

Selection of Subjects

Seven of the nine studies enrolled subjects from selected hospitals. No enumerated underlying population census exists from which to draw a sample of the study base population, so various approximations are employed. In Cramer et al (1982) and Harlow et al (1992), controls were drawn from the towns in which the cases reside. This is a reasonable approach that assumes women from the same towns reflect the medical referral population from which the cases

CONFIDENTIAL

were admitted to the hospitals. The overall relative risk for use of powders after bathing in these two studies were 1.9 and 1.5. The relative risks associated with the known risk factors in these two studies were generally similar with estimates from all other case-control studies and from cohort studies.

In Hartge et al (1983), Whittemore et al (1988), Booth et al (1989) and Rosenblatt et al (1992), controls were drawn from women admitted to the same hospitals as cases for conditions unrelated to talc or any of the known or suspected risk factors under investigation. This also is a sound approach often applied in studies with cases from selected hospitals without an enumerated underlying population. The relative risks from these studies ranged from 0.7 to 2.4, but the two studies with substantial numbers and some details on exposure (Booth et al, 1989 and Whittemore et al, 1988) were closer to each other with relative risks of 1.3 and 1.4, respectively.

The study by Tzonou selected controls from people who were visiting the cases. This is a less common approach that may yield a control group more like the case group in social class and other variables than is the underlying study base. On the other hand, the protective effect of parity and oral contraceptive use was seen in this study. Nevertheless, the relative risk due to talc exposure was 0.9.

Cases from an enumerated population were used in the studies by Harlow and Weiss (1989) in Seattle (Relative risk = 1.1) and by Chen et al (1992) in Beijing, China (RR = 3.9). The former study was limited to borderline ovarian cancers, but the etiologic similarity of these and invasive cancers has been amply demonstrated (Harlow et al, 1988).

CONFIDENTIAL

Thus, the basic design of these nine case-control studies shows no clear pattern of effects being seen with one design but not others. There may be a very slight gradient of decreasing risk estimates going from hospital-town studies (highest), to hospital-hospital studies, to the one U.S. population-based study. This pattern was also observed in the extensive analysis of population-based studies and hospital-based studies by Whittemore et al (1992), which showed consistent but subtle effects of study design in twelve case-control studies of ovarian cancer. For example, the population-based studies showed a slightly stronger effect of oral contraceptive use than did the hospital-based studies, but the protection by oral contraceptives was readily apparent in both (Whittemore et al, 1992). The differences reflect the combination of two distinct effects: hospitalized patients are heavier, generally less healthy, slightly poorer and otherwise not perfectly representative of the general population-based cases and controls (a difference that favors using population-based cases and controls whenever possible); but hospitalized patients generally are more willing to participate in studies (a difference that may make hospital-based studies superior depending upon response rates).

Participation Rates

In all the U.S. studies, participation rates were similar to each other and in the low but acceptable range, 69%-77% in cases and 60%-80% in controls. Rates were not available from the study in Britain (Booth et al, 1989) but were likely to be higher than in the United States. Response rates were very high, in excess of 90%, in China and Greece, but talc use was so uncommon in these regions that estimates of the talc effect were quite unstable (as measured in the confidence intervals). Thus, differences in participation rates among the studies are not likely to explain differences in talc findings.

In general, a high rate of non-response in either cases or controls can distort results, and a much higher rate in one or the other causes alarm. Neither of these problems appear to be present in the nine studies. There also is no reason to suspect that talc use is moderately or strongly related to willingness to participate. While bias from non-response cannot be utterly discounted, it does not appear to be an important or obvious problem in these data.

Recollection of Talc Exposures

If exposures to talc were recalled differently by cases and controls, relative risk estimates would be distorted. Cases could over-report talc use if they were anxious to assign a cause to their cancer, or underreport it if they were anxious to avoid providing this information. However, this differential response bias would have to be claimed to occur for talc and not for the numerous other exposures queried in these interviews that typically last for one hour and involve more than 100 questions. Differential misclassification, and random error, are most likely to occur if the question is too long or too vague, and if the behavior itself is intermittent and trivial.

In the nine studies, the questions pertaining to talc were generally short. The behavior is trivial and not memorable, but habitual for many and thus not particularly likely to be misremembered. Forthcoming data from the Nurses Health Study (Hankinson et al, 1993), a prospective cohort analysis, will relate ovarian cancer to talc habits queried before diagnosis. Until then, recollection bias should not be regarded as a likely explanation for the talc effect.

Analysis of Data

Three features of analysis should be considered: a) uncorrected confounding effects of factors that cause ovarian cancer and are related to talc use, b) analysis of the data to separate distinct aspects of the exposure, particularly dose, route, type and timing, and c) analysis of the combined effect of talc and other factors. None of the studies show obvious failure to control for the effects of potential confounders. Data from Harlow et al (1992) illustrate that talc use does vary with age, parity and oral contraceptive use. These effects are controlled in all of the overall relative risk estimates under review. How plausible is it that unidentified confounders are at work? If weight is a confounder, as suggested by Rosenblatt et al (1992), then it must have a stronger relation to ovarian cancer risk than has been reported to date. If the association with talc is actually null, then a confounder needs to be very closely linked to talc and must have an effect on ovarian cancer risk that is higher than the 1.4-1.8 estimated for talc.

Source of exposure does show a pattern in the studies reviewed, with direct application to the perineum carrying the greatest risk. Brands do not appear to be separable. Dose-response data are available from only a few studies (Tables 2-3). Frequency shows an effect in two of the three studies, and duration in one of the two. For duration and cumulative dose, the relative risk rises to 1.8, and possibly higher in women whose use occurs during ovulatory time periods. The typical user applied talc daily for at least a decade.

Possible co-factors include tubal ligation, hysterectomy, tubal occlusions, parity and sexual activity because they might affect the likelihood of talc

migration to the ovary, and indicators of ovulation which may affect ovarian cancer susceptibility. Various techniques are employed to assess these types of interactions.

We have elaborated on the existing data reported in the study by Harlow et al (1992) to illustrate the joint effect of talc and number of ovulations (Table 4). The effect of talc is stronger among women with fewer than the median number of ovulations. It is possible that the talc effect is most easily seen against a background of lower risk. On the other hand, the data suggest that the effects of talc and ovulation do not exasperbate each other. However, this simple analysis does not separate talc exposure occurring during ovulatory and nonovulatory time periods. When applications of talc during nonovulatory time periods were evaluated using data from Harlow et al (1992), there was no excess ovarian cancer risk associated with increasing applications of talc during nonovulatory time periods (data not shown), and virtually all of the ovarian cancer risk occurred as a consequence of talc exposure during time periods when women were ovulatory. We can therefore not rule out the possibility that this pattern may represent an interaction with ovulation.

Table 4 also shows the joint effect of talc and mid-cycle pain - a symptom of ovulation. Although talc does appear to operate most strongly in women reporting mid-cycle pain, such pain is not related to risk, suggesting that it is not a reliable index of ovulation. Lastly, the joint effect of talc and gynecologic surgery in data from Whittemore et al (1988) is also shown in Table 4. The association with talc does not appear to depend on whether tubal ligation or hysterectomy had occurred. It is certainly possible that talc exerts an effect in the years before surgery. A more detailed analysis of timing might uncover such an effect, but a large study would be required.

Since Cramer et al (1982) presented epidemiologic data in support of the hypothesis that exposure to cosmetic talc is associated with increased risk of developing ovarian cancer, eight additional case-control studies have provided data to assess the potential association. Of these, four also have implicated use of body powders, three with odds ratios of 1.5 or more. If talcum powder use cannot induce ovarian cancer, what is likely to explain the positive studies? The play of chance is possible, but this is not where suspicion most logically points. The cumulative effect of several modest errors during data collection, perhaps not even the same errors in all studies, theoretically could produce these findings. However, no obvious error in collection or analysis clearly explain the studies with positive findings, and one must regard the range of relative risk estimates from epidemiology, 1.0 to 1.8, as plausible.

Additional epidemiologic studies, especially those with different designs, may help, but the greatest need is to confirm or deny the reports of talc embedded in human ovarian tissue and the report of easy transportation of particles through the human female reproductive tract.

- Acheson, E.D., Gardner, M.J., Pippard, E.C., and Grime, L.P. (1982). Mortality of two groups of women who manufactured gas masks from chrysotile and crocidolite asbestos: a 40-year follow-up. Br. J. Ind. Med. 39 (4), 344-348.
- Booth, M., Beral, V., and Smith, P. (1989). Risk factors for ovarian cancer: A case control study. Br. J. Cancer. 60 (4), 592-598.
- Chen, Y., Wu, P., Lang, Y., Ge, W., Hartge, P., and Brinton, L.A. (1992). Risk factors for epithelial ovarian cancer in Beijing, China. Int. J. Epidemiol. 21 (1), 23-29.
- C.T.F.A. Specification (1976). Talc, cosmetic: Cosmetic, toiletry and fragrance association. Inc. Issue 1976, 10-17.
- Cralley, L.J., Key, M.M., Groth, D.H., et al (1968). Fibrous and mineral content of cosmetic talcum products. Am. Ind. Hyg. Assoc. J. 29 (4), 350-354.
- Blejer, H.P., Arlon, R. (1973). Talc: A possible occupational and environmental carcinogen. J. Occup. Med. 15 (2), 92-97.
- Cramer, D.W., Welch, W.R., Scully, R.E., and Wojciechowski, C.A. (1982). Ovarian cancer and talc. Cancer. 50 (2), 372-376.

- Egli, G.E. and Newton, M. (1961). The transport of carbon particles in the human female reproductive tract. Fertil Steril. 12, 151-155.
- Eiseman, B., Seelig, M.G., and Womack, N.A. (1947). Talcum powder granuloma: a frequent and serious postoperative complication. Ann. Surg. 126 (5), 820-832.
- Graham, J. and Graham, R. (1967). Ovarian cancer and asbestos. Environ. Res. 1 (2), 115-128.
- Hankinson, S.E., Hunter, D.J., Colditz, G.A., Willett, W.C., Stampfer, M.J., Rosner, B., Hennekens, C.H., Speizer, F.E. (1993). J Am Med Assoc. 270 (23), 2813-2818.
- Harlow, B.L., Weiss, N.S., Roth, G.J., Chu, J., and Daling, J.R. (1988). Case-control study of borderline ovarian tumors: Reproductive history and exposure to exogenous female hormones. Cancer Res. 48 (20), 5849-5852.
- Harlow, B.L. and Weiss, N.S. (1989). A case-control study of borderline ovarian tumors: the influence of perineal exposure to talc. Am. J. Epidemiol. 130 (2), 390-394.
- Harlow, B.L., Cramer, D.W., Bell, D.A., and Welch, W.R. (1992). Perineal exposure to talc and ovarian cancer risk. Obstet. Gynecol. 80 (1), 19-26.
- Hartge, P., Hoover, R., Leshner, L.P., and McGowan, L. (1983). Talc and ovarian cancer (letter). J. Am. Med. Assoc. 250 (14), 1844.

- Henderson, W.J., Joslin, C.A.F., Turnbull, A.C., and Griffiths, K. (1971). Talc and carcinoma of the ovary and cervix. J. Obstet. Gynecol. Brit. Common. 78 (3), 266-272.
- Henderson, W.J., Hamilton, T.C., and Griffiths, K. (1979). Talc in normal and malignant ovarian tissue [Letter]. Lancet. 1 (8114), 499.
- Henderson, W.J., Hamilton, T.C., Baylis, M.S., Pierrepont, C.G., and Griffiths, K. (1986). The demonstration of the migration of talc from the vagina and posterior uterus to the ovary in the rat. Environ. Res. 40 (2), 247-250.
- Henderson, W.J. (1969). A simple replication technique for the study of biological tissues by electron microscopy. J. Microscopy. 89 (3), 369-372.
- Hildick-Smith, G.Y. (1976). The biology of talc. Br. J. Ind. Med. 33 (4), 217-229.
- Keal, E.E. (1960) Asbestosis and abdominal neoplasms. Lancet. 2, 1211-1216.
- Kleinfeld, M., Messite, J., and Zaki, M.H. (1974). Mortality experience among talc workers: a followup study. J. Occup. Med. 16 (5), 345-349.
- Longo, D.L., and Young, R.C. (1979). Cosmetic talc and ovarian cancer [Letter]. Lancet. 2 (8150), 1011-1012.
- Michaels, L., and Chissick, S.S. (1979). Asbestos, properties, applications and hazards volume 1. (ed), pp 1-5, John Wiley & Sons Ltd,

- Newhouse, M.L., Berry, G., and Wagner, J.C. (1985) Mortality of factory workers in east London 1933-80. Br. J. Ind. Med. 42 (1), 4-11.
- Parnley, T.H., and Woodruff, J.D. (1974) The ovarian mesothelioma. Am. J. Obstet. Gynecol. 120 (2), 234-241.
- Rohl, A.N., Langer, A.M., Selikoff, I.J., et al (1976). Consumer talcums and powders: mineral and chemical characterization. J. Toxicol. Environ. Health. 2 (2), 255-284.
- Rosenblatt, K.A., Szklo, M., and Rosenshein, N.B. (1992). Mineral fiber exposure and the development of ovarian cancer. Gynecol. Oncol. 45 (1), 20-25.
- Tobacman, J.K., Greene, M.H., Tucker, M.A., et al (1982) Intra-abdominal carcinomatosis after prophylactic oophorectomy in ovarian cancer-prone families. Lancet. 2 (8302), 795-797;
- Thomas, T.L., and Stewart, P.A. (1987) Mortality from lung cancer and respiratory disease among pottery workers exposed to silica and talc. Am J. Epidemiol. 14 (5), 35-43.
- Tzonou, A., Polychronopoulou, A., Hsieh, C., Rebelakos, A., Karakatsani, A., and Trichopoulos, D. (1993). Hair dyes, analgesics, tranquilizers and perineal talc application as risk factors for ovarian cancer. Int. J. Cancer. 55 (3), 408-410.

- Venter, P.F. and Iturralde, M. (1979). Migration of particulate radioactive tracer from the vagina to the peritoneal cavity and ovaries. S. Afr. Med. J. 55 (23), 917-919.
- Whittemore, A.S., Wu, M.L., Paffenbarger, R.S., Sarles, D.L., Kambert, J.B., Grosser, S., Jung, D.L., Ballon, S., and Hendrickson, M. (1988). Personal and environmental characteristics related to epithelial ovarian cancer. II. Exposures to talcum powder, tobacco, alcohol, and coffee. Am. J. Epidemiol. 128 (6), 1228-1240.
- Whittemore, A.S., Harris, R., and Itnyre, J. (1992). Characteristics relating to ovarian cancer risk: Collaborative analysis of 12 U.S. case-control studies. II. Invasive epithelial ovarian cancers in white women. Collaborative ovarian cancer group. Am J Epidemiol 136(10), 1184-1203.
- Wignall, B.K., and Fox, A.J. (1982). Mortality of female gas mask assemblers. Br. J. Ind. Med. 39 (1), 34-38.

Table 1. Relative risk estimates (with 95% confidence intervals) of ovarian cancer in relation to perineal exposure to talc

First Author	(Year)	Any Genital Exposure	Application to Sanitary Napkins	Diaphragm Exposure	Direct application after bathing
Cramer	(1982)	1.9 (1.3-2.9)	1.5 (0.7-3.2)	1.6 (0.6-3.9)	1.6 (1.0-2.7)
Hartge	(1983)	2.5 (0.7-10.0)	b	0.8 (0.4-1.4)	b
Whittemore	(1988)	1.4 (0.9-1.8)	0.6 (0.2-1.8)	1.5 (0.6-3.6)	1.5 (0.8-2.6)
Harlow	(1989) ^a	1.1 (0.7-2.1)	1.9 (0.9-6.9)	0.5 (0.2-1.3)	1.2 (0.6-2.6)
Booth	(1989)	1.3 (0.9-1.9)	b	0.7 (0.4-1.1)	b
Harlow	(1992)	1.5 (0.9-1.8)	1.1 (0.4-2.8)	1.2 (0.6-2.4)	1.7 (1.1-2.7)
Chen	(1992)	3.9 (0.9-10.6)	b	b	b
Rosenblatt	(1992)	2.4 (1.1-5.3)	4.8 (1.3-17.8)	3.0 (0.8-10.8)	1.7 (0.7-3.9)
Tzonou	(1993)	0.9 (0.3-2.7)	b	b	b

^aRestricted to borderline ovarian tumors.

^bData not available.

Table 2. Relative risk estimates (with 95% confidence intervals) of ovarian cancer in relation to frequency and years of exposure to talc

	Frequency of Exposure			Years of Exposure	
	None ^a	<Daily	Daily	<10	>10
Whittemore (1988)	1.0	1.3 (0.8-2.0)	1.5 (0.9-2.2)	1.6 (1.0-2.6)	1.1 (0.7-1.7)
Booth (1989)	1.0	1.4 (0.9-2.1)	1.3 (0.8-1.9)	b	b
Harlow (1992)	1.0	1.4 (0.7-2.4)	1.8 (1.1-3.0)	1.2 (0.5-2.6)	1.6 (1.0-2.7)

^aReferent group.

^bData not available

Table 3. Relative risk estimates^a (with 95% confidence intervals) of ovarian cancer in relation to lifetime applications of perineal talc exposure

	Lifetime Applications			
	None	<1000	1000-10,000	>10,000
All exposures	1.0	1.3 (0.7-2.7)	1.5 (0.9-2.4)	1.8 (1.0-3.0)
Exposure prior to tubal ligation or hysterectomy only	1.0	1.4 (0.7-2.9)	1.5 (0.9-2.4)	1.7 (1.0-3.0)
Exposure prior to tubal ligation or hysterectomy, and during ovulatory periods only	1.0	1.5 (0.8-2.9)	1.3 (0.8-2.0)	2.8 (1.4-5.4)

^aAdjusted for parity (0, 1-2, ≥3), education (<12 years, ≥12 years), marital status (never married, ever married), religion (jewish, nonjewish), use of sanitary napkins (no, yes), douching (no, yes), age (continuous), and weight (<140lbs, ≥140lbs).

Table 4. Relative risk estimates* of ovarian cancer in relation to perineal talc exposure by control median months of ovulation, history of midcycle pain or gynecologic surgery.

		Number of Ovulations ^a		Midcycle Pain ^a		Gynecological Surgery ^b	
		<Median	≥Median	No Pain	Any Pain	No Surg.	Surgery
No	Cases:Controls	49:80	72:65	93:107	28:38	70:182	21:110
Talc	Odds Ratios ^c	1.0 ^d	2.2	1.0 ^d	0.9	1.0 ^d	0.5
Any	Cases:Controls	40:39	74:55	88:77	26:17	71:151	26:96
Talc	Odds Ratios ^c	2.1	3.3	1.4	1.7	1.3	0.8

^aData from Harlow et al, 1992.

^bData from Whittemore et al, 1988.

^cAdjusted for parity (0, 1-2, ≥3), education (<12 years, ≥12 years), marital status (never married, ever married), religion (jewish, nonjewish), use of sanitary napkins (no, yes), douching (no, yes), age (continuous), and weight (<140lbs, ≥140lbs) for odds ratios stratified by ovulations and midcycle pain, and only parity for odds ratios stratified by gynecological surgery.

^dReferent group